

TOPOLOGICALLY NONTRIVIAL SECTORS OF MAXWELL FIELD THEORY ON RIEMANN SURFACES

FRANCO FERRARI

*Dipartimento di Fisica, Università di Trento, 38050 Povo, Italy
and INFN, Gruppo Collegato di Trento*

ABSTRACT

In this paper the Maxwell field theory is considered on a closed and orientable Riemann surface of genus $h > 1$. The solutions of the Maxwell equations corresponding to nontrivial values of the first Chern class are explicitly constructed for any metric in terms of the prime form.

August 1996

Two dimensional gauge field theories on manifolds have recently been considered by various authors [1]–[13]. In this paper the Maxwell field theory is treated on a closed and orientable Riemann surface of genus $h > 1$. The explicit construction of the propagators and of the flat connections has been already discussed for the more general Yang–Mills field theories in ref. [14]. However, also the abelian case is interesting because of the presence of topologically nontrivial sectors. For instance, the topologically nontrivial solutions of the Maxwell equations have applications both in string theory [15] and in the Fractional Quantum Hall Effect [16]. On Riemann surfaces equipped with arbitrary metrics a rigorous and explicit derivation of the gauge connections labelled by nonzero values of the first Chern class [17] is still missing in the physical literature. This derivation is the main result of this letter. To this purpose, we first compute the general solutions of the Maxwell field theory corresponding to a total magnetic flux Φ in terms of the prime form [18] and of the $2h$ holomorphic differentials. For general values of Φ , these solutions are periodic around the homology cycles of the Riemann surface. However, we show here that the periodicity amounts to a well defined gauge transformation if Φ satisfies the Dirac quantization condition. With our formulas it is possible for example to explicitly verify the fact that the wave functions of the Landau levels in the Quantum Hall effect are independent of the metric as indirectly proved in [16].

Let us start from the action of the Maxwell field theory which is given by:

$$S_{Maxwell} = \int_M dx^1 \wedge dx^2 \frac{\sqrt{|g(x)|}}{4} F_{\mu\nu} F^{\mu\nu} \quad (1)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ and M is a general closed and orientable Riemann surface of genus $h > 1$ parametrized by the real coordinates x^μ , $\mu = 1, 2$. From now on we will put $dx^1 \wedge dx^2 \equiv d^2x$ for simplicity. On M we define an Euclidean metric $g_{\mu\nu}(x)$ with determinant $g(x) = \det g_{\mu\nu}$ and a canonical basis of homology cycles a_i and b_i , with $i = 1, \dots, h$. Let us now choose local complex coordinates $z = x^1 + ix^2$ and $\bar{z} = x^1 - ix^2$. In these variables the determinant of the metric $g_{\alpha\beta}$, $\alpha, \beta = z, \bar{z}$ is given by $g(z, \bar{z}) = \frac{1}{4}g(x)$. Explicitly, $g(z, \bar{z}) = g_{zz}g_{\bar{z}\bar{z}} - g_{z\bar{z}}g_{\bar{z}z}$. The h holomorphic differentials admitted on M will be denoted with the symbols $\omega_i(z)dz$, $i = 1, \dots, h$. We assume that they are normalized in such a way that $\oint_{a_i} dz \omega_j(z) = \delta_{ij}$. As a consequence, the matrix

$$\Omega_{ij} \equiv \oint_{b_i} \omega_j(z)dz$$

is the so-called period matrix. For future purposes, it will also be necessary to recall the basics of the Hodge decomposition on Riemann surfaces without boundary. Any vector field $(A_z, A_{\bar{z}})$ can be decomposed according to the Hodge decomposition theorem in coexact, exact and harmonic components as follows:

$$A_z = \partial_z \phi + \partial_z \rho + A_z^{\text{har}} \quad A_{\bar{z}} = -\partial_{\bar{z}} \phi + \partial_{\bar{z}} \rho + A_{\bar{z}}^{\text{har}} \quad (2)$$

The coexact and exact components are expressed using the two scalar fields ϕ and ρ respectively. Here ϕ is purely complex, while ρ is real. A_z^{har} and $A_{\bar{z}}^{\text{har}}$ take into account of the presence of the holomorphic differentials. The decomposition (2) is not invertible unless

$$\int_M d^2 z \sqrt{|g(z, \bar{z})|} \phi(z, \bar{z}) = \int_M d^2 z \sqrt{|g(z, \bar{z})|} \rho(z, \bar{z}) = 0 \quad (3)$$

Here $d^2 z \sqrt{g} \equiv i \sqrt{g} d\bar{z} \wedge dz$ is the volume form on M in complex coordinates. Let us now compute the equations of motion coming from the action (1). Noting that in two dimensions the only non-vanishing components of the field strength $F_{\mu\nu}$ are

$$F_{z\bar{z}} = -F_{\bar{z}z} = \partial_z A_{\bar{z}} - \partial_{\bar{z}} A_z \quad (4)$$

an easy computation shows that the gauge fields satisfy the following equations of motion:

$$\partial_{\bar{z}} \left[\frac{1}{\sqrt{|g(z, \bar{z})|}} (\partial_{\bar{z}} A_z - \partial_z A_{\bar{z}}) \right] = 0 \quad (5)$$

$$\partial_z \left[\frac{1}{\sqrt{|g(z, \bar{z})|}} (\partial_z A_{\bar{z}} - \partial_{\bar{z}} A_z) \right] = 0 \quad (6)$$

The solutions of the above equations are of the form [4], [16]:

$$F_{z\bar{z}} dz \wedge d\bar{z} = \frac{2\pi i \Phi \sqrt{|g(z, \bar{z})|}}{A} d\bar{z} \wedge dz \quad (7)$$

where $A = \int_M d^2 x \sqrt{|g(x)|}$ is the area of the surface and Φ is a real constant representing the total magnetic flux. To show this is a simple exercise, but it will be useful to fix the notation. Let us perform the change of variables $(z, \bar{z}) \rightarrow (x^1, x^2)$ in the integral $I = \int_M d^2 z F_{z\bar{z}}$. Since $d^2 z = -2i d^2 x$ and $F_{12} = -2i F_{z\bar{z}}$, where $F_{12} = \partial_1 A_2 - \partial_2 A_1$, one finds that $I = \int_M d^2 x F_{12}$. Thus I is proportional to the first Chern class defined by:

$$k = \frac{1}{2\pi} \int_M d^2 x F_{12} \quad k = 0, \pm 1, \dots \quad (8)$$

On the other side, from eq. (7), we have that $I = 2\pi i\Phi \int_M d^2z \sqrt{|g(z, \bar{z})|} \frac{d\bar{z} \wedge dz}{A}$. Remembering that $2\sqrt{g(z, \bar{z})} = \sqrt{g(x^1, x^2)}$, we find the relation $d^2z \sqrt{|g(z, \bar{z})|} = d^2x \sqrt{g(x^1, x^2)}$. As a consequence $I = 2\pi\Phi$ and thus we obtain the desired relation giving Φ as the total magnetic flux in or outgoing from the Riemann surface:

$$2\pi\Phi = \int_M d^2x F_{12} \quad (9)$$

Because of the Dirac quantization of the flux, the topologically non-trivial solutions of the equations of motion (5)–(6) are characterized by the following integer values of Φ :

$$\Phi = k \quad k = 0, \pm 1, \pm 2, \dots \quad (10)$$

Eq. (7) is equivalent to the following set of equations:

$$\partial_z A_{\bar{z}} = \frac{i\pi C_1 \sqrt{|g(z, \bar{z})|}}{A} \quad \partial_{\bar{z}} A_z = \frac{i\pi C_2 \sqrt{|g(z, \bar{z})|}}{A} \quad (11)$$

provided the fields A_z and $A_{\bar{z}}$ are coexact forms. For consistency with eq. (9), the two constants C_1 and C_2 must obey the following relation: $C_1 - C_2 = 2\Phi$. To solve eq. (11) explicitly in the gauge fields, we introduce the following Green function [19]:

$$K(z, w) = \log|E(z, w)|^2 - 2\pi \sum_{i,j=1}^h \left[\text{Im} \int_w^z \omega_j(s) ds \right] (\text{Im } \Omega)_{ij}^{-1} \left[\text{Im} \int_w^z \omega_j(s) ds \right] \quad (12)$$

where $\text{Im } a = \frac{a - \bar{a}}{2i}$ denotes the imaginary part of a complex number a . $E(z, w)$ is the prime form [18] and $|\text{Im } \Omega|_{ij}^{-1}$ is the inverse of the matrix $\frac{(\Omega - \bar{\Omega})_{ij}}{2i}$. $K(z, w)$ satisfies the equation:

$$\partial_z \partial_{\bar{z}} K(z, w) = \pi \delta_{z\bar{z}}^{(2)}(z, w) - \pi \sum_{i,j=1}^h \omega_i(z) (\text{Im } \Omega)_{ij}^{-1} \bar{\omega}_j(\bar{z}) \quad (13)$$

We will show now that a particular solution of eqs. (11), uniquely defined modulo holomorphic connections, is:

$$A_z^I = \frac{iC_1}{A} \int_M d^2w \partial_z K(z, w) \sqrt{|g(w, \bar{w})|} + i\pi C_1 \sum_{i,j=1}^h \omega_i(z) (\text{Im } \Omega)_{ij}^{-1} \int_{\bar{z}_0}^{\bar{z}} \bar{\omega}_j(\bar{w}) d\bar{w} \quad (14)$$

$$A_{\bar{z}}^I = \frac{iC_2}{A} \int_M d^2w \partial_{\bar{z}} K(z, w) \sqrt{|g(w, \bar{w})|} + i\pi C_2 \sum_{i,j=1}^h \bar{\omega}_i(\bar{z}) (\text{Im } \Omega)_{ij}^{-1} \int_{z_0}^z \omega_j(w) dw \quad (15)$$

First of all, it is easy to see with the help of eq. (13) that the above gauge fields really satisfy eqs. (11). Moreover, since the field strength (4) is completely antisymmetric in its indices, it takes no contributions from exact components. Indeed, the above vector fields A_z^I and $A_{\bar{z}}^I$ are purely coexact 1-forms. To verify this, we note that A_z^I and $A_{\bar{z}}^I$ can be obtained, apart from a constant factor, after derivation of the purely complex function

$$\phi^I(z, \bar{z}) = \frac{i}{A} \int_M d^2w \sqrt{|g(w, \bar{w})|} K(z, w) + i\pi \sum_{i,j=1}^h \int_{z_0}^z \omega_i(w) dw (\text{Im } \Omega)_{ij}^{-1} \int_{\bar{z}_0}^{\bar{z}} \bar{\omega}_j(\bar{w}) d\bar{w} \quad (16)$$

Exploiting the symmetry properties of the period matrix Ω in its indices, we have in fact:

$$A_z^I = C_1 \partial_z \phi^I \quad A_{\bar{z}}^I = C_2 \partial_{\bar{z}} \phi^I \quad (17)$$

Comparing with eq. (2), it is easy to see that the gauge fields A_α^I of eqs. (14)–(15) are coexact only if $C_1 = -C_2$. This condition, together with eq. (8), fixes the constants C_1 and C_2 uniquely:

$$C_1 = -C_2 = \Phi \quad (18)$$

Finally we have to check that the particular solutions A_z^I and $A_{\bar{z}}^I$ given in (14)–(15) remain single-valued, apart from a well defined gauge transformation, when transported around any nontrivial homology cycle γ . Indeed, from their definitions (14)–(15), it turns out that the gauge fields A_z^I and $A_{\bar{z}}^I$ are periodic around γ :

$$A_z'^I = A_z^I + iC_1\pi \sum_{i,j=1}^h \omega_i(z) (\text{Im } \Omega)_{ij} \oint_{\gamma} \bar{\omega}_j(\bar{w}) d\bar{w} \quad (19)$$

$$A_{\bar{z}}'^I = A_{\bar{z}}^I + iC_2\pi \sum_{i,j=1}^h \bar{\omega}_i(\bar{z}) (\text{Im } \Omega)_{ij} \oint_{\gamma} \omega_j(w) dw \quad (20)$$

These transformations amount however to a $U(1)$ gauge transformation

$$A_\alpha'^I = A_\alpha^I + iU^{-1}(z, \bar{z}) \partial_\alpha U(z, \bar{z}) \quad \alpha = z, \bar{z} \quad (21)$$

where the element of the gauge group is provided by:

$$U(z, \bar{z}) = U(z_0, \bar{z}_0) \exp \left[\pi C_1 \sum_{i,j=1}^h \left(\int_{z_0}^z \omega_i(s) ds (\text{Im } \Omega)_{ij} \oint_{\gamma} \bar{\omega}_j(\bar{w}) d\bar{w} - \right. \right.$$

$$\int_{\bar{z}_0}^{\bar{z}} \bar{\omega}_i(\bar{s}) d\bar{s} (\text{Im}\Omega)_{ij} \oint_{\gamma} \omega_j(w) dw \Big) \Big] \quad (22)$$

First of all, substituting the above defined group element in eq. (21), one obtains the transformations (19)–(20) as expected. Moreover, it is easy to check that the exponent appearing in the right hand side of eq. (22) is purely imaginary. As a consequence it turns out that $\overline{U(z, \bar{z})} = U^{-1}(z, \bar{z})$. However, this is still not enough to conclude that $U(z, \bar{z})$ is really an element of $U(1)$, because one should also verify that $U(z, \bar{z})$ is not multivalued when transported around a nontrivial homology cycle. To this purpose, let us first consider the case in which $\gamma = b_l$, with $l = 1, \dots, h$. After a slight reshuffling of indices, we have from eq. (22):

$$U(z, \bar{z})|_{\gamma=b_l} = U(z_0, \bar{z}_0)|_{\gamma=b_l} \exp \left[2\pi i C_1 \sum_{i,j=1}^h \left(\bar{\Omega}_{li} (\Omega - \bar{\Omega})_{ij}^{-1} \int_{z_0}^z \omega_i(s) ds - \Omega_{li} (\Omega - \bar{\Omega})_{ij}^{-1} \int_{\bar{z}_0}^{\bar{z}} \bar{\omega}_i(\bar{s}) d\bar{s} \right) \right]$$

At this point we recall that the real harmonic differentials α_l , defined in such a way that

$$\oint_{a_k} \alpha_l = \delta_{kl} \quad \oint_{b_k} \alpha_l = 0 \quad (23)$$

have the following expression [20]:

$$\alpha_l = \sum_{i,j=1}^h (-\bar{\Omega}_{li} (\Omega - \bar{\Omega})_{ij}^{-1} \omega(z) dz + c.c.)$$

where *c.c.* stands for complex conjugate. Therefore, setting $z_0 = (x_0^1, x_0^2)$ and $z = (x^1, x^2)$, $U(z, \bar{z})$ becomes in real coordinates:

$$U(x)|_{\gamma=b_l} = U(x_0)|_{\gamma=b_l} \exp \left[-2\pi i C_1 \int_{(x_0^1, x_0^2)}^{(x^1, x^2)} \alpha_l \right]$$

Here we have used the relation $\int_{\sigma} dx^{\mu} A_{\mu} = \int_{\sigma} (dz A_z + d\bar{z} A_{\bar{z}})$, valid for any vector field A_{μ} and for any path σ on M . Now we move $U(x)$ along a closed cycle b_k . Let us denote the result of this operation with $U'(x)$. Since the exponent in the right hand side of eq. (22) is purely imaginary, $U'(x)$ should differ from $U(x)$ at most by a complex phase. However, exploiting eqs. (23), it is easy to see that:

$$U'(x) = U(x)$$

Analogously, transporting $U(x)$ around a cycle a_k , we have:

$$U'(x) = U(x)e^{-2\pi i C_1 \delta_{kl}}$$

Remembering the condition (10) on the magnetic flux and eq. (18), it is clear that $e^{-2\pi i C_1 \delta_{kl}} = 1$, proving that $U(z, \bar{z})$ is singlevalued around the b -cycles.

To conclude our proof, we consider the case in which $\gamma = a_l$ in eq. (22). Performing similar calculations as before, one finds:

$$U(x)|_{\gamma=a_l} = U(x_0)|_{\gamma=a_l} \exp \left[2\pi i C_1 \int_{(x_0^1, x_0^2)}^{(x^1, x^2)} \beta_l \right]$$

where the symbols β_l , $l = 1, \dots, h$, denote now the real harmonic differentials

$$\beta_l = \sum_{i=1}^h [(\Omega - \bar{\Omega})_{li}^{-1} \omega_i(z) dz - (\Omega - \bar{\Omega})_{li}^{-1} \bar{\omega}_i(\bar{z}) d\bar{z}]$$

They are normalized along the homology cycles as follows [20]:

$$\oint_{a_k} \beta_l = 0 \qquad \oint_{b_k} \beta_l = \delta_{kl}$$

Exploiting the above equations, a straightforward calculation shows that $U(x)$ is singlevalued also when transported across an a -cycle. This completes our proof that $U(z, \bar{z})$ is a well defined element of the group $U(1)$. As a consequence, the claim has been verified, that the gauge fields given in (14)–(15) are particular solutions of the Maxwell equations (5)–(6) on any closed and orientable Riemann surface M equipped with an arbitrary metric $g_{\mu\nu}$. These solutions correspond to topologically non-trivial configurations with nonvanishing first Chern class. If the total flux Φ is not an integer, the periodicity of A_α^I around the homology cycles, computed in eqs. (19)–(20), cannot be reabsorbed by a gauge transformation, since $U(z, \bar{z})$ starts to be multivalued and thus it is no longer a valid group element on M . All the other solutions corresponding to nonvanishing values of the first Chern class, can be obtained from eqs. (14)–(15) adding linear combinations of the harmonic differentials and exact forms.

Concluding, we have computed here the nontrivial solutions of the Maxwell equations. The Dirac quantization of the total flux Φ naturally arises from the requirement that the periodicity of these solutions around the homology cycles can be reabsorbed by a well defined gauge transformation. As briefly mentioned in the Introduction, there are some applications of our results both in the quantum Hall effect and in the theory of superstrings with $N = 2$. These topics lay however beyond the scope of this short letter and will be treated elsewhere.

References

- [1] M. Atiyah and R. Bott, *Phil. Trans. R. Soc. Lond.* **A 308** (1982), 523.
- [2] E. Witten, *Comm. Math. Phys.* **141** (1991), 153; *Jour. of Geom. and Phys.* **9** (1992), 3781.
- [3] D. J. Gross *Nucl. Phys.* **B400** (1993), 161; D. J. Gross and Washington Taylor IV, *Nucl. Phys.* **B400** (1993), 181; *ibid.* **B403** (1993), 395
- [4] G. Thompson, *1992 Trieste Lectures on Topological Gauge Theory and Yang–Mills Theory*, Preprint IC/93/112 (hep-th/9305120).
- [5] A. Bassetto, L. Griguolo and G. Nardelli, *Yang–Mills Theories on the Space–Time $S_1 \times \mathbf{R}$: Equal Time Quantization in Light Cone Gauge and Wilson Loops*, Preprint DFPD-95-TH-07 (hep-th/9506095); A. Bassetto and L. Griguolo, *Nucl. Phys.* **B439** (1995), 327.
- [6] K. S. Gupta, R. J. Henderson. S. G. Rajeev and O. T. Torgut, *Jour. Math. Phys.* **35** (1994), 3845; S. G. Rajeev, *Phys. Lett.* **212B** (1988), 203.
- [7] I. Sachs and A. Wipf, *Helv. Phys. Acta* **65** (1992), 653; *Phys. Lett* **326B** (1994), 105.
- [8] E. Abdalla, M.C.B. Abdalla and K. D. Rothe, *Nonperturbative Methods in Two Dimensional Quantum Field Theory*, Singapore 1991, World Scientific.
- [9] N. J. Hitchin, *Gauge Theories on Riemann Surfaces*, in Lectures on Riemann Surfaces, M. Cornalba & al. (eds), ICTP Trieste Italy, 9 Nov.-18 Dec. 1987, World Scientific editions; *Topology*, **31** (3) (1992), 449; *Proc. London Math. Soc.* **55**, 59 (1987).
- [10] D. S. Fine, *Comm. Math. Phys.* **134** (1990), 273; **140** (1991), 321; *Quantum Yang–Mills on a Riemann Surface II: Topological Sectors and Measures on Moduli Space*, (hep-th 9504103).
- [11] A. Sengupta, *Comm. Math. Phys.* **147** (1992), 191; *Ann. Phys.* **221** (1993), 17.
- [12] H. Joos, *Nucl. phys.* **B17** (Proc. Suppl.) (1990), 704; *Helv. Phys. Acta* **63** (1990), 670.
- [13] C. Jayewardena, *Helv. Phys. Acta* **61** (1988), 636.
- [14] F. Ferrari, *Helv. Phys. Acta* **67** (1994), 702.
- [15] S. V. Ketov and O. Lechtenfeld, *Phys. Lett.* **353B** (1995), 463.
- [16] R. Iengo and Dingping Li, *Nucl. Phys.* **B413** (1994), 735.
- [17] T. Eguchi, P. B. Gilkey and A. J. Hanson, *Phys. Rep.* **66** (1980), 213.
- [18] J. D. Fay, *Lect. Notes in Math.* **352**, Springer Verlag, 1973.
- [19] E. Verlinde and H. Verlinde, *Nucl. Phys.* **B288** (1987), 357.
- [20] H. M. Farkas and I. Kra, *Riemann Surfaces*, Springer Verlag, Berlin, Hedelberg, New York 1980.